

Knowledge, Attitudes, and Practices about Trachoma in Four Evaluation Units of Low Endemic Areas, in Yemen: Implications for Prevention and Control

Sami A. Al-Haidari^{1*}, Mansour A. Al-Taj¹, Abdulkareem A. Nassar¹, Yahia A. Raja'a¹, & Tawfik Al-Khateeb²

¹Department of Community Medicine, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen ²Department of Ophthalmology, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen

Corresponding Author: Sami A. Al-Haidari, Department of Community Medicine, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen. Orchid ID: https://orcid.org/0000-0002-4706-0778

Cite: Al-Haidari SA, Al-Taj MA, Nassar AA, Raja'a YA, Al-Khateeb T. Knowledge, attitudes, and practices about trachoma in four evaluation units of low endemic areas, in Yemen: Implications for prevention and control. Cryst J Med Res. 2025;1(1): 01-12.

Received: March 04, 2025; Accepted: March 15, 2025; Published: 21 March, 2025

Abstract

Background: Trachoma is a bacterial infection that causes blindness worldwide and affects impoverished communities disproportionately. Despite these interventions, Yemen has made strides towards trachoma elimination.

Objective: To assess the knowledge, attitudes, and practices (KAP) regarding trachoma and its associated factors among rural communities in six districts within four Evaluation Units (EUs) of Hodeidah and Ibb governorates in, Yemen.

Methods: A community-based cross-sectional study was conducted in six districts from four Evaluation Unites (EUs) of the Hodeidah and Ibb governorates in January 2025. Bivariate analysis (chi-square) identified variables for multivariable logistic regression (p < 0.25). Multicollinearity was addressed by excluding relevant variables from each model.

Results: Among the 1164 interviewed participants, 38.8% (435) exhibited good knowledge, 19.1% (222) demonstrated good attitudes, and 46.5% (541) practiced good hygiene related to trachoma prevention. Significant associations were observed between KAP and several factors: Age of the head of household: Inversely associated with adequate knowledge (AOR = 0.13, 95% CI 0.08-0.22, P < 0.001), good attitudes (AOR = 3.12, 95% CI 2.23-4.36, P < 0.001), and good practices (AOR = 0.26, 95% CI 0.18-0.39, P < 0.001). Mother's literacy status: Significantly and positively associated with adequate knowledge (AOR = 102, 95% CI 47.7-219.4, P < 0.001), good attitudes (AOR = 3.60, 95% CI 2.50-5.17, P < 0.001), and good practices (AOR = 170, 95% CI 22.2-85.5, P < 0.001) and good practices (AOR = 3.67, 95% CI 2.35-5.73, P < 0.001), but inversely associated with good attitudes (AOR = 0.43, 95% CI 0.31-0.61, P < 0.001). Household size: Inversely associated with adequate knowledge (AOR = 0.62, 95% CI 0.40-0.99, P < 0.001) and good attitudes (AOR = 1.82, 95% CI 1.33-2.50, P < 0.001). Wealth index: Significantly and positively associated with adequate with adequate knowledge (AOR = 0.62, 95% CI 0.40-0.99, P < 0.001) and good attitudes (AOR = 1.82, 95% CI 3.94-10.62, P < 0.001) and good practices (AOR = 1.57, 95% CI 1.03-2.40, P < 0.001). Gender of the head of household: Male-headed households were less likely to have a good knowledge than female-headed households (AOR = 0.59, 95% CI 0.37-0.96, P < 0.001).

Conclusions: This study found significant disparities in KAP related to trachoma prevention, influenced by factors like the age of the household head, parental literacy, and household wealth.

Keywords

Trachoma, Evaluation Unites, KAP

Author Summaries

Trachoma, a bacterial infection causing blindness, disproportionately affects impoverished communities. While Yemen has made progress toward eliminating this disease, it remains a public health concern. This study explored community knowledge, attitudes, and practices (KAP) regarding trachoma in rural areas of Hodeidah and Ibb governorates. Researchers interviewed over 1,1164 people and found that while some had good hygiene practices (46.5%), knowledge (38.8%) and positive attitudes (19.1%) about trachoma prevention were less common. The study revealed strong links between KAP and factors like the age and gender of the head of household, parental literacy, family size, and wealth. For example, children whose mothers were literate were significantly more likely to have good trachoma-related knowledge, attitudes, and practices. Similarly, wealthier households tended to have better understanding and practices These findings highlight the need for targeted interventions that address these social and economic disparities to effectively combat trachoma in Yemen.

Introduction

Trachoma, a bacterial infection of the eye, constitutes the second leading cause of preventable blindness worldwide. This treatable and preventable disease disproportionately affects impoverished populations. (1-4) Trachoma disproportionately affects individuals in disadvantaged rural areas with limited access to necessities like clean water and sanitation. (5, 6) Globally, an estimated 142.2 million people are at risk of trachoma, and it is responsible for visual impairment in approximately 1.9 million individuals. (7, 8) The World Health Organization (WHO) endorses the SAFE (Surgery, Antibiotics, Facial cleanliness, and Environmental improvement) strategy for trachoma elimination, which has been implemented in over 40 countries worldwide where trachoma is endemic. (9, 10) The SAFE strategy can eliminate trachoma, but its proper application remains crucial as the disease continues to pose a global health threat. (11). Trachoma persists as a significant health issue in the world's poorest communities. Several factors contribute to an increased risk of trachoma, including limited access to clean water, poor personal hygiene, and inadequate environmental sanitation. (12-14). In

Yemen 2024, the adjusted prevalence of the clinical sign trachomatous inflammation-follicular (15) among children aged 1-9 years in the enumeration units encompassing highly endemic areas specifically Hodeidah and Ibb was 0.9 % and 0.8%, respectively.) Unpublished (Trachoma is a substantial segment of the population that exhibits unfavorable attitudes and a lack of knowledge concerning trachoma infection. Although the implementation of various interventions, including annual mass drug administration (MDA), for trachoma control has expanded in Yemen in recent years, significant knowledge gaps persist among vulnerable populations, particularly those residing in remote rural areas and those experiencing economic hardship.(16) These populations may have limited awareness and adherence to recommended trachoma prevention and control measures. However, there is currently a dearth of data on KAP regarding trachoma prevention and control within rural communities in the six districts. The research findings shed light on critical gaps in knowledge, community practices, and perceptions surrounding trachoma, and how these factors influence patient health-seeking behavior. This valuable information is crucial for promoting health and preventing trachoma by addressing the underlying causes and drivers of disease spread. Ultimately, these findings can significantly enhance the impact of programs aimed at preventing and eliminating trachoma within the country. Furthermore, these findings will provide valuable insights for policymakers in formulating appropriate interventions to implement the SAFE strategy effectively. However, the study aimed to assess the KAP to give us more concepts for eliminating the disease.

Methods

Study Design, Population and Setting

A community-based cross-sectional study was conducted among households in six low-endemic trachoma districts across four Evaluation Units (EUs) in January 2025. These districts, identified in 2024 with a TF prevalence exceeding 5%, included Far Aludayan and Mudaykhirah in Ibb Governorate, and Almighalf, Almunirah, As Salif, and Azayadyah in Hodeidah Governorate (Fig. 1). Eligible participants were male and female heads of households who provided informed consent.



Figure 1: Prevalence of trachomatous inflammation (TF) < 5% in four evaluion unite (EU) Hodaida and Ibb

Sample Size Calculation

The sample size was calculated to be 1164 using standard epidemiological procedures. This calculation incorporated the following assumptions: a 95% confidence level ($Z\alpha/2 = 1.96$), an estimated population proportion of 60% with adequate knowledge regarding trachoma, a desired margin of error of 5%, a design effect of 2.63, and an inflation factor of 1.2 to account for potential data loss. (17)

Sampling Strategy

A two-stage cluster sampling approach was utilized. In the first stage, 30 clusters were systematically selected from villages within the study districts, and grouped into (EUs) according to WHO criteria. (18) In the second stage, a compact segment sampling method was used to select households from each cluster, with selection proportional to cluster population size. Households were defined as individuals residing in a dwelling unit, regardless of marital status. Households were then randomly selected from the lists. Informed consent was obtained from the heads of selected households (or their adult representatives) before study participation, as per WHO guidelines. (18)

Data Collection

A structured questionnaire was adapted from a previous study. (19, 20) Face-to-face interviews with eligible household heads were conducted. Direct observations were used to measure personal and household practices. A qualitative method was used to determine the attitudes of the community toward the trachoma. Before conducting interviews with respondents, the team was trained on research tools. To prevent disease transmission among participants, investigators used disposable gloves or cleaned their hands with 70% alcohol.

Operational Definition

Knowledge was assessed using nine questions, with correct answers scored as 1 and incorrect as 0. Good knowledge was defined as scoring above the median score. Attitudes and practices were similarly assessed using seven questions each. Individuals scoring above the median on the attitude and practice scales were categorized as having good attitudes, and good practices, respectively. Therefore, participants scoring above the median of 7, 7, and 24 were considered to have good knowledge, practice, and attitude, respectively.

Statistical Analysis

SPSS version 26 was used for data analysis. All the total KAP score variables were not normally distributed, as determined by the Shapiro-Wilk test (p-value ≤ 0.05). However, the median was used as the cutoff point for the categorizing of the total knowledge, practice, and attitude scores. Frequency, and percentage were used to present categorical variables.

Bivariate analyses were conducted to identify potential associations between independent and dependent variables. Variables with p-values less than 0.25 in the bivariate analyses were included in a multivariable logistic regression model. Multicollinearity was checked using variance inflation factors (VIF), and variables with high multicollinearity were removed. For example, the practice variable was removed during the multivariate analysis for knowledge, the knowledge variable was removed during the multivariate analysis for practice, and both knowledge and practice variables were removed during the multivariate analysis for attitude. A p-value < 0.05 was considered statistically significant.

Results

Characteristics of the Study Population

A majority of household heads were female (68.3%), and aged 40 years or younger (57.6%), and a substantial portion of households had more than 7 members (49.1%). Most fathers were literate (79.0%) and mothers were literate (58.7%), and the majority of fathers were employed (77.2%). Also, a significant proportion of households were classified as "poor" in terms of the Wealth index (38.8%) (Table 1).

 Table 1: Sociodemographic characteristics of study participants (N=1164).

Characteristics	n	(%)
Gender of the household head		
Male	369	31.7
Female	795	68.3
Age of the household head (years)		
<u>≤</u> 40	671	57.6
>40	493	42.4
Family size (members)		
≤7	592	50.9
>7	572	49.1
Father's literacy status		
Literate	919	79.0
Illiterate	245	21.0
Mother's literacy status		
Literate	683	58.7
Illiterate	481	41.3

Father's employment status		
Employed	899	77.2
Unemployed	265	22.8
Wealth index		
Rich	712	61.2
Poor	452	38.8

Knowledge of Trachoma

A majority 983 (84.5%) of the study subjects had heard about trachoma. However, 722 (62.0%) of them had heard through MDA. Two hundred and thirty-seven (20.4%) of the respondents correctly answered the cause of trachoma. Regarding measures that can be taken to prevent trachoma 402 (34.5%) daily hand washing with water and soap, symptoms of redness and tearing of the eye 352 (30.2%),362 (31.1%) respectively. Transmission

of trachoma, 570 (46.6%), 90 (7.7%), and 175 (15.0%) of the study subjects correctly responded that contact with an unhygienic environment, through flies landing on the face and contact with nasal or eye discharges respectively. 941(80.5%) knowing trachoma can be treated. Less than two quarters, 435 (38.8%) of the study subjects had a good knowledge of trachoma (Table 2).

Characteristics	n	(%)
Heard about trachoma disease	899	77.2
Yes	983	84.5
No	181	15.5
By what have you heard	712	61.2
MDA	722	62.0
Community	90	7.7
TV	56	4.8
Radio	98	8.4
Others	198	17.0
What do you think causes trachoma		
Bacteria	237	20.4
Dust	542	46.6
Flies	201	17.3
Virus	74	6.4
I don't know	110	9.5
What measures can be taken to prevent trachoma		
Avoid contact with nasal or eye discharges	88	7.6
Avoid sharing clothes	125	10.7
Daily hand wash with water and soap	402	34.5
Improving access to clean water	141	12.1
Use medicine	408	35.1
What are the symptoms of trachoma that you know		
Itching	219	18.8
Photophobia	86	7.4
Redness of the eye	352	30.2
Scratching of the eyelashes	145	12.5
Tearing of the eye	362	31.1
How is trachoma transmitted?		
Unhygienic environment	570	46.6
Through flies landing on the face	90	7.7
Contact with nasal or eye discharges	175	15.0
Sharing clothes or fomites	206	17.7

I don't know	110	9.5
How often should a person wash their face to prevent trachoma		
One	157	13.5
Twice	185	15.9
Three times	452	38.8
When it dirty	370	31.8
Can trachoma be treated?		
Yes	941	80.8
No	155	13.3
I don't know	68	5.8
Where would you go if a family member had symptoms of trachoma		
Hospital	351	30.2
Local Clinic	298	25.6
Ophthalmologist	207	17.8
Traditional Healer	217	18.6
I don't know	91	7.8
Knowledge status		
Good	435	38.8
Poor	712	61.2

Attitude on Trachoma Prevention and Control

Nine hundred and forty-two (80.9%) of the study participants had a poor attitude toward trachoma infection. According to this study, 471 (40.5%) strongly agreed that trachoma is a health problem in the community. Conversely, 535 (46.0%) agreed and 426 (36.6%) agreed that Flies are the main vector of the disease and transmitted from person to person. 369 (31.7%) of the

study subjects disagreed that trachoma may eventually lead to blindness. Nearly 370 (31.8%),334 (28.7%), and 458 (39.3%) of the respondents agreed and strongly agreed that hygiene and sanitation are preventing disease, disposing of waste is a protection from disease, and taking drugs is important to prevent and control, respectively (Table 3).

Table 3: Attitude of the Study Community Toward Trachoma in Four Evaluation Units in Yemen, 2025.

Variables	Likert Scale					
	Agree	Strongly agree	Disagree	Strongly disagree	Neutral	
Trachoma is a health problem in your community	392 (33.7%)	471(40.5%)	123(10.6%)	127(10.9%)	51(4.4%)	
Flies are the main vector of the disease	535(46.0%)	219(18.8%)	142(12.2%)	101(8.7%)	167(14.3%)	
Trachoma transmitted from person to person	426(36.6%)	273(23.5%)	168(14.4%)	147(12.6%)	150(12.9%)	
The disease may eventually lead to blindness.	132(11.3%)	145(12.5%)	191(16.4%)	369(31.7%)	327(28.1%)	
Hygiene and sanitation are a preventing disease	370(31.8%)	288(24.7%)	301(25.9%)	183(15.7%)	22(1.9%)	
Disposing of waste is a protection from disease	332(28.5%)	334(28.7%)	103(8.8%)	291(25.0%)	104(8.9%)	
Taking drugs is important to prevent and control	458(39.3%)	360(30.9%)	96(8.2%)	106(9.1%)	144(12.4%)	
Attitude Category						
Good			222 (19.1%)			
Poor	942 (80.9%)					

Practices on Trachoma Prevention and Control

The majority of respondents, 348 (29.9%), and 347(29.7%) reported that the main source of water for domestic use during the dry season was from wells and springs respectively. 862 (74.1%) reported a distance between home and water source of less than 30 minutes. The majority,1003 (86.2%) reported no presence of flies on the child's face and 987 (84.8%) reported no presence of

animals in the house. However, 781 (67.1%),667 (57.3%) and 912 (78.4%) used enough water for bathing, had proper solid and liquid waste disposal management, and utilized a latrine, respectively. Out of the total, 541 (46.5%) respondents were classified as having good practices towards trachoma prevention and control (Table 4).

Characteristics	n	(%)
The main source of water for domestic use during the dry season		
Rainwater	118	10.1
River	87	7.5
Spring	347	29.8
Transportation tankers	264	22.7
Well	348	29.9
Distance between home and water source	941	80.8
<30 Min	862	74.1
>30Min	302	25.9
The presence of flies on the child's face		
No	1003	86.2
Yes	161	13.8
The presence the feces around the house		
No	987	84.8
Yes	177	15.2
The presence of animals in the house		
No	541	46.5
Yes	623	53.5
Use enough water for bathing		
No	383	32.9
Yes	781	67.1
Having proper solid &liquid waste disposal management		
No	497	42.7
Yes	667	57.3
Latrine utilization		
No	252	21.6
Yes	912	78.4
Practice Category		
Good	541	46.5
Poor	623	53.5

Table 4: Practices of study Community of trachoma in Four Evaluation Units in Yemen, 2025.

Multivariable Logistic Regression Analysis for Factors Associated with Knowledge of Study Participants about Trachoma in Four Evaluation Units in Yemen, 2025

Multivariable logistic regression analysis revealed that age of the head of household, gender of the head of household, household size categories, mother's literacy status, father's occupation status, and wealth index were significantly associated with knowledge status of the participants AOR=0.13,95% CI:0.08–0.22), (AOR=0.59,95% CI:0.37–0.96), (AOR=0.62,95% CI:0.40– 0.99), (AOR=102,95% CI:47.7–219.4), (AOR=43.6,95%-CI:22.2–85.5), and AOR=6.47, 95% CI: 3.94–10.62), respectively (Table 5).

Table 5: Logistic regression analysis of selected variables with knowledge of study participants on trachoma in Four Evaluation

 Units in Yemen, 2025.

Variables	Knowledge		COR (95% CI)	P-value	AOR (95% CI)	P- value	
	Adequate %	Inadequate %					
Age of the head of household (years)							
≤40	75(16.6%)	418(58.7%)	0.14(0.11-0.19)	< 0.001	0.13(0.08-0.22)	< 0.001	
>40	377(83.4%)	294(41.3%)	Ref		Ref		
Gender of the head of household							
Male	96(21.2%)	273(38.3%)	0.43(0.33-0.57)	< 0.001	0.59(0.37-0.96)	0.033	

Female	356(78.8%)	439(61.7%)	Ref		Ref		
Household size categories							
≤7	195(43.1%)	397(55.8%)	0.60(0.48-0.76)	< 0.001	0.62(0.40-0.99)	0.044	
>7	257(56.9%)	315(44.2%)	Ref		Ref		
		F	Father's literacy status				
Literate	373(98.2%)	239(33.6%)	1.44(1.065 - 1.935)	0.01	0.59(0.33-1.06)	0.076	
Illiterate	8(1.8%)	473(66.4%)	Ref		Ref		
		Ν	Iother's literacy status				
Literate	444(86.6%)	82(17.4%)	109(53.7-224.8)	< 0.001	102(47.7-219.4)	< 0.001	
Illiterate	93(13.4%)	388(82.6%)	Ref		Ref		
		Fa	ther's occupation status	S			
Employed	438(96.9%)	461(64.7%)	17.03(9.79-29.64)	< 0.001	43.6(22.2-85.5)	< 0.001	
Unemployed	14(3.1%)	251(35.3%)	Ref		Ref		
			Wealth Index				
Rich	366(81.0%)	346(48.6%)	4.50(3.41-5.94)	< 0.001	6.47(3.94-10.62)	< 0.001	
Poor	86(19.0%)	366(51.4%)	Ref		Ref		
			Practice Category				
Good	451(100.0%)	89(12.5%)	6.08(5.03-7.35)	< 0.001			
Poor	0(0.0%)	623(87.5%)	Ref				
			Attitude Category				
Good	66(14.6%)	156(21.9%)	0.61(0.44-0.84)	0.001	0.98(0.57-1.69)	0.939	
Poor	386(85.4%)	556(78.1%)	Ref		Ref		

Multivariable Logistic Regression Analysis for Factors Associated with Attitudes of Study Participants about Trachoma in Four Evaluation Units in Yemen, 2025

Multivariable logistic regression analysis showed that age of the head of household, household size categories, Mother's literacy

status, and father's occupation status, were significantly associated with attitudes status of the participants AOR=3.12, 95%CI: 2.23–4.36), AOR=1.82, 95%CI: 1.33–2.50), AOR=3.60, 95%CI: 2.50–5.17) and AOR=0.43, 95%CI: 0.31–0.61) respectively (Table 6).

Table 6: Logistic regression analysis of selected variables with Attitude of study participants on trachoma in Four Evaluation Units in Yemen, 2025.

Variables	Attitudes		COR (95% CI)	P-value	AOR (95% CI)	P- value			
	Good (%)	Poor (%)							
	Age of the head of household (years)								
≤40	123(55.4%)	370(39.3%)	1.92(1.43-2.58)	< 0.001	3.12(2.23-4.36)	< 0.001			
>40	99(44.6%)	572(60.7%)	Ref		Ref				
		Gende	r of the head of house	chold					
Male	148(66.7%)	647(68.7%)	0.91(0.67-1.25)	0.307					
Female	74(33.3%)	295(31.3%)	Ref						
		Но	usehold size categorie	S					
≤7	135(60.8%)	457(48.5%)	1.65(1.22-2.22)	0.001	1.82(1.33-2.50)	< 0.001			
>7	87(39.2%)	485(51.5%)	Ref		Ref				
		F	ather's literacy status						
Literate	45(20.3%)	200(21.2%)	0.94(0.656-1.36)	0.415					
Illiterate	177(79.7%)	742(78.8%)	Ref						
		M	Iother's literacy status						
Literate	160(72.1%)	523(55.5%)	2.07(1.50-2.85)	< 0.001	3.60(2.50-5.17)	< 0.001			
Illiterate	62(27.9%)	419(44.5%)	Ref		Ref				
		Fat	her's occupation statu	IS					
Employed	147(66.2%)	752(79.8%)	0.50(0.36-0.68)	< 0.001	0.43(0.31-0.61)	< 0.001			

Cryst Jour Medic Res 2025

Unemployed	75(38.8%)	190(20.2%)	Ref		Ref				
	Family ownership of land								
Rich	133(59.9%)	579(61.5%)	0.94(0.70-1.26)	0.362					
Poor	89(40.1%)	363(38.5%)	Ref						
		ŀ	Knowledge Category						
Good	66(29.7%)	386(41.0%)	0.61(0.44-0.84)	0.001					
Poor	156(70.3%)	556(59.0%)	Ref						
Practice Category									
Good	155(69.8%)	386(41.0%)	3.33(2.43-4.56)	< 0.001					
Poor	67(30.2%)	556(59.0%)	Ref						

Multivariable Logistic Regression Analysis for Factors Associated with Practices of Study Participants about Trachoma in Four Evaluation Units in Yemen, 2025

pation status, and wealth index were significantly associated with practices status of the participants AOR=0.26, 95%CI: 0.18–0.39), AOR=170, 95%CI: 82–358), AOR=3.67, 95%CI: 2.35–5.73), and AOR=1.57, 95%CI: 1.03–2.40), respectively (Table 7).

Multivariable logistic regression analysis indicated that age of the head of household, mother's literacy status, father's occu-

 Table 7: Logistic regression analysis of selected variables with Practice of study participants on trachoma in Four Evaluation Units in Yemen, 2025.

Variables	Prac	tices	COR (95% CI)	P-value	AOR (95% CI)	P- value		
	Good (%)	Poor (%)						
		Age of t	he head of household ((years)				
≤40	106(19.6%)	387(62.1%)	0.15(0.11-0.19)	< 0.001	0.26(0.18-0.39)	< 0.001		
>40	435(80.4%)	236(37.9%)	Ref		Ref			
Gender of the head of household								
Male	133(24.6%)	236(37.9%)	0.54(0.42-0.69)	< 0.001	0.70(0.47-1.06)	0.088		
Female	408(75.4%)	387(62.1%)	Ref		Ref			
		Но	usehold size categorie	S				
≤7	234(43.3%)	358(57.5%)	0.56(0.45-0.71)	< 0.001	0.75(0.50-1.11)	0.148		
>7	307(56.7%)	265(42.5%)	Ref		Ref			
		F	ather's literacy status					
Literate	105(19.4%)	140(22.5%)	1.20(0.91-1.60)	0.114				
Illiterate	436(80.6%)	483(77.5%)	Ref					
		N	Iother's literacy status					
Literate	533(98.5%)	150(24.1%)	210(102-432)	< 0.001	170(82-358)	< 0.001		
Illiterate	8(1.5%)	473(75.9%)	Ref		Ref			
		Fat	ther's occupation statu	S				
Employed	457(84.5%)	442(70.9%)	2.23(1.67-2.98)	< 0.001	3.67(2.35-5.73)	< 0.001		
Unemployed	84(15.5%)	181(29.1%)	Ref		Ref			
			Wealth Index					
Rich	380(70.2%)	332(53.3%)	2.07(1.62-2.65)	< 0.001	1.57(1.03-2.40)	0.038		
Poor	161(29.8%)	291(46.7%)	Ref		Ref			
	Knowledge Category							
Good	452(83.5%)	623(100.0%)	0.12(0.10-0.15)	< 0.001				
Poor	89(16.5%)	0(0.0%)	Ref					
			Attitude Category					
Good	155(28.7%)	67(10.8%)	3.33(2.43-4.56)	< 0.001				
Poor	386(71.3%)	556(89.2%)	Ref					

Discussion

The study revealed significant associations between good knowledge, attitudes, and practices regarding trachoma and several factors, including the age of the head of household, the mother's literacy status, and the father's occupation status. Good knowledge and good attitudes were significantly associated with household size, while good knowledge and good practices were significantly associated with the wealth index. Additionally, the gender of the head of the household was significantly associated with good knowledge.

Regarding knowledge, the majority of respondents (84.5%) had heard of trachoma, comparable to findings in Northern Ethiopia (89.2%) and Bangladesh (86%) (19, 21). However, this rate was higher than that reported in Kenya (65.7%) and lower than in other countries (22, 23), suggesting that community mobilization and sensitization efforts may have been implemented in the study sites. MDA campaigns were the primary source of information for 62% of participants, suggesting that these initiatives have significantly contributed to raising awareness about trachoma. Similar to studies in other countries like Zambia, Ethiopia, and Kenya. (24-26)The present study revealed that (46.6%), (20.4%), and (17.3%) of respondents reported correctly that trachoma can be transmitted by dust, bacteria, and flies respectively, Similar studies from Kenya and Ethiopia reported that the most reported mode of trachoma transmission was contact with flies and dirt. (21, 27) and daily handwashing with soap and water and the use of medication were identified as the most common preventive measures. While less frequent, other preventive measures mentioned, such as avoiding contact with eye discharges and sharing clothes, demonstrate some understanding of potential trachoma transmission routes. This suggests a gap in the understanding of trachoma's true etiology, and the likely reason would be the lack of awareness of the participants toward trachoma this finding is in line with a study done in other countries. (3, 28, 29)Participants identified several potential transmission routes for trachoma, including unhygienic environments, contact with nasal or eye discharges, and the sharing of clothes or contaminated items. They also recognized that trachoma can be prevented by going to the hospital or health facility, seeing specialists, use of appropriate medications, improving sanitation, and regular hand and face washing with soap. Similar findings have been reported in studies conducted in other countries. (30-32)Finally, (61.2%) of the community demonstrated inadequate knowledge regarding trachoma.

Regarding attitudes, a majority (74.2%) of respondents agreed or strongly agreed that trachoma is a health problem in their community, indicating a general awareness of the issue. Similar findings have been reported in studies conducted in other countries. (20, 28, 30, 33)A considerable proportion (64.8%) agreed or strongly agreed that flies are the main vector of the disease, highlighting a significant misconception about trachoma transmission, (59.1%) agreed or strongly agreed that trachoma is transmitted from person to person, indicating a partial understanding of transmission routes. (34, 35)(23.8%) agreed or strongly agreed that trachoma may eventually lead to blindness, suggesting limited awareness of the potential severity of the disease. (36-38)A majority (56.5%) agreed or strongly agreed that hygiene and sanitation are important for preventing the disease, indicating some understanding of preventive measures. A large proportion (70.2%) agreed or strongly agreed that taking drugs

is important for prevention, and control, suggesting a positive perception of medical interventions. Overall Attitude: A significant proportion (80.9%) of the community demonstrated poor attitudes towards trachoma, characterized by misconceptions about transmission, limited awareness of potential consequences, and inadequate understanding of preventive measures.it is crucial to emphasize that medication is part of a comprehensive approach to trachoma control and should be used in conjunction with other preventive measures. (14, 39-41) Regarding practices, water source, and accessibility: the majority of the households (74.1%) had access to water sources within a 30-minute walk, Springs (29.8%) and wells (29.9%) were the common sources of domestic water. (42-44) Hygiene Practices: A significant proportion of households (67.1%) reported using enough water for bathing A significant proportion of households (86.2%) reported no presence of flies on children's faces. (45, 46) Environmental Factors: The majority of households (84.8%) reported no presence of feces around the house. More than half of the households (53.5%) reported the presence of animals inside the house. (33, 47, 48) Reducing the burden of flies and improving environmental sanitation through proper human excreta disposal are crucial for public health. (49) Waste Disposal: A substantial proportion of households (57.3%) reported having proper solid and liquid waste disposal management. Latrine utilization was high, with 78.4% of households reporting using a latrine. (50, 51) Overall Practices: A significant proportion (53.5%) of households demonstrated poor practices related to trachoma prevention. (19, 28).

Regarding reparation multivariate knowledge, the study found that adequate trachoma knowledge was linked to several socioeconomic factors, including the age of the head of the household, parental literacy, father's occupation, and household wealth. This suggests that socioeconomic disparities significantly influence knowledge levels within the community. The finding that female-headed households demonstrated higher levels of adequate knowledge is a significant observation and warrants further investigation. (52) The strong association between good practices and adequate knowledge highlights the importance of promoting and reinforcing positive health behaviors within the community. Targeted interventions are needed to address knowledge gaps among vulnerable populations, particularly those from low-income households and those with limited access to education. (53, 54).

Regarding reparation multivariate attitudes, Several socioeconomic factors, such as the age of the head of household, household size, mother's literacy status, and father's occupation status, were significantly associated with attitudes towards trachoma. (32, 44) This suggests that socioeconomic disparities play a crucial role in shaping community attitudes. The finding that households with good knowledge were less likely to have good attitudes towards trachoma is unexpected and warrants further investigation. It is possible that individuals with some knowledge may be more aware of the complexities of the disease and may have developed more nuanced (and potentially more negative) perspectives. Improving community knowledge about trachoma is crucial for fostering positive attitudes and promoting effective preventive behaviors. Interventions to improve attitudes towards trachoma should be tailored to address the specific needs and concerns of different population groups, such as younger heads of households and those from low-income households.

Regarding reparation multivariate practices, several socioeconomic factors, such as the age of the head of household, the mother's literacy status, the father's occupation status, and household wealth, were significantly associated with good practices related to trachoma prevention. This emphasizes the importance of addressing socioeconomic disparities to improve health outcomes. The strong association between good knowledge and good practices highlights the crucial role of health education in promoting positive health behaviors. The finding that maternal literacy is significantly associated with good practices underscores the importance of women's education in improving household health outcomes. Trachoma prevention interventions should be tailored to address the specific needs and circumstances of different population groups, such as low-income households and those with limited access to education.

Conclusion and Recommendation

Our study suggests that even in areas with low trachoma prevalence, community KAP regarding trachoma prevention and control must be enhanced to achieve elimination. This necessitates health education campaigns focused on the SAFE strategy, implemented at the community level, during mass drug administration campaigns, and at health facilities. The aim is to increase knowledge, address cultural perceptions that hinder behavior change, and ultimately promote positive behaviors for trachoma prevention and control within communities. Interestingly, our findings revealed higher levels of knowledge among female-headed households, warranting further investigation into the factors contributing to this observation. Notably, we observed an association between good knowledge and less favorable attitudes toward trachoma prevention and control, suggesting a need for further research to understand this unexpected relationship and tailor interventions accordingly.

Acknowledgments

The authors thank the head household and their families for participation in the study. They also thank the ophthalmologists who examined the children. Special thanks are due to the team from the districts and governorates targeted.

Author Contributions

SAA conceived the study idea and implemented the fieldwork. SAA, MAA, and AHN designed the study. SAA, MAA, and AHN analyzed the data, interpreted the results, and drafted the manuscript. All authors read, revised, and approved the final version of the manuscript.

Funding

The authors received no financial support for the research, authorship, and/ or publication of this article.

Availability of Data and Materials

All relevant data are included in the manuscript, and the datasets for the study are available from the corresponding author upon reasonable request.

Declarations

Ethics Approval and Consent to Participate

The protocol of this study was approved by the Research Ethics Committee of the Faculty of Medicine and Health Sciences at Sana'a University, Yemen. Oral informed consent was obtained from the heads of the households or their legal representatives after they were briefed about the nature and purpose of the study. The privacy of participants and the confidentiality of their data were assured. The study was conducted according to the guidelines of the Declaration of Helsinki.

Consent for Publication

Not applicable.

Competing Interests

The authors declare that they have no competing interests.

References

- Gross LL. Oromo Ethiopians Perceptions of the Prevalence, Causes, Treatment and Prevention of Trachoma: Walden University; 2019.
- 2. Okoye R. Exploring public health intervention strategies to address barriers and challenges in the prevention of avoidable blindness due to glaucoma in Anambra State, Nigeria: Middlesex University; 2020.
- Kamau JW. Prevalence, Risk Factors and Trachoma Causing Species Circulating in East Pokot, Baringo County, Kenya 2023.
- Gebre T, Kello AB, Habtamu E, Ngondi JM. Trachoma. Neglected Tropical Diseases-Sub-Saharan Africa: Springer; 2024. p. 415-31.
- 5. Gupta V, Gupta N, Senjam S, Vashist PJNTD-SA. Trachoma. 2017:219-44.
- 6. Shafi Abdurahman O, Last A, Macleod D, Habtamu E, Versteeg B, Dumessa G, et al. Trachoma risk factors in Oromia Region, Ethiopia. 2023;17(11):e0011679.
- Yafanna G, Musa D, Abubakar N, Abubakar H, Kode S, Alhassan MJBJoN, et al. Awareness of Trachoma Prevention among People of Bolori Community Maiduguri Metropolitan Council Local Government Area of Borno State, Nigeria. 2022;4(1):949-58.
- 8. Kurian M, Nozad BJFJotNSfMB. How is the World Health Organization (WHO) tackling neglected tropical diseases affecting the eye? The case of trachoma and onchocerciasis. 2023;6(1):12-29.
- Organization WH. Report of the 21st meeting of the WHO alliance for the global elimination of trachoma by 2020, Geneva, Switzerland, 20-22 April 2017. World Health Organization; 2019.
- Renneker KK, Abdala M, Addy J, Al-Khatib T, Amer K, Badiane MD, et al. Global progress toward the elimination of active trachoma: an analysis of 38 countries. 2022;10(4):e491-e500.
- Lavett DK, Lansingh VC, Carter MJ, Eckert KA, Silva JC-JTSWJ. Will the SAFE strategy be sufficient to eliminate trachoma by 2020? Puzzlements and possible solutions. 2013;2013(1):648106.
- 12. Abebe TA, Tucho GTJPNTD. The impact of access to water supply and sanitation on the prevalence of active trachoma in Ethiopia: A systematic review and meta-analysis. 2021;15(9):e0009644.
- 13. Garn JV, Boisson S, Willis R, Bakhtiari A, Al-Khatib T, Amer K, et al. Sanitation and water supply coverage thresholds associated with active trachoma: modeling cross-sectional data from 13 countries. 2018;12(1):e0006110.
- 14. Ageed A, Khan MJC. Eliminating Trachoma in Africa: The Importance of Environmental Interventions. 2024;16(1).

- 15. Uzoka F-M, Akwaowo C, Nwafor-Okoli C, Ekpin V, Nwokoro C, El Hussein M, et al. Risk factors for some tropical diseases in an African country. 2021;21(1):2261.
- 16. Alhaidari SA, Raja'a YA, Mahdy MA, Haider TA, Nassar A. Knowledge, Attitude, Practices and Associated Factors Towards Trachoma Among Communities in Highly Endemic Areas, Yemen. Sana'a University Journal of Medicine and Health Sciences. 2025;19(1):135-43.
- 17. Mengistu K, Shegaze M, Woldemichael K, Gesesew H, Markos YJCO. Prevalence and factors associated with trachoma among children aged 1–9 years in Zala district, Gamo Gofa Zone, Southern Ethiopia. 2016:1663-70.
- Organization WH. Design parameters for population-based trachoma prevalence surveys: strategic and technical advisory group for neglected tropical diseases, working group on monitoring and evaluation. World Health Organization; 2018.
- Gebretnsae H, Mamo N, Teklemariam T, Fenta K, Gebrehiwet T, Berhe A, et al. Knowledge, attitudes, and practices about trachoma in rural communities of Tigray Region, Northern Ethiopia: implications for prevention and control. 2020;2020(1):3270530.
- 20. Churko C, Asfaw MA, Zerdo ZJCO. Knowledge, attitude, practices and associated factors towards trachoma among people living in Arba Minch Zuria District, Gamo Zone, Southern Ethiopia. 2021:3075-85.
- 21. Belaynew W, Berihun M, Tadesse A, Yared AJJoO. Knowledge and practice on childhood blindness among communities in Northwest Ethiopia: implications to blindness prevention programs. 2013.
- 22. Islam FMA, Chakrabarti R, Islam SZ, Finger RP, Critchley CJPO. Factors associated with awareness, attitudes and practices regarding common eye diseases in the general population in a rural district in Bangladesh: the Bangladesh population-based diabetes and eye study (BPDES). 2015;10(7):e0133043.
- Ng'etich AS, Owino C, Juma A, Khisa KNJJophiA. Knowledge, attitudes and eye care seeking practices regarding trachoma in Central Division of Kajiado County, Kenya. 2015;6(1).
- 24. Maritim CP. Evaluation of implementation determinants shaping the appropriateness of a trachoma mass drug administration program in Livingstone district, Zambia: The University of Zambia; 2017.
- 25. Astale T, Sata E, Zerihun M, Nute AW, Stewart AE, Gessese D, et al. Population-based coverage survey results following the mass drug administration of azithromycin for the treatment of trachoma in Amhara, Ethiopia. 2018;12(2):e0006270.
- 26. Gichuki PM, Kimani BW, Kanyui T, Okoyo C, Watitu T, Omondi WP, et al. Using community-based participatory approaches to improve access to mass drug administration for trachoma elimination in a pastoral conflict area of Kenya. 2024;18(11):e0012653.
- 27. Munguti P, Ng'ang'a Z, Muttunga JJE, Journal CAM. Knowledge, practices and perception on trachoma and its influence on health seeking behaviour of the pastoralist patients in Kajiado Central Division. 2015;2:89-96.
- 28. Njomo DW, Karimurio J, Odhiambo GO, Mukuria M, Wanyama EB, Rono HK, et al. Knowledge, practices and perceptions of trachoma and its control among communities of Narok County, Kenya. 2016;2:1-10.

- 29. MWANIKI S. Factors affecting trachoma control and prevention among pastoralist in Kirisia sub-county, Samburu county, Kenya: Maseno university; 2024.
- Thompson K, Hutchins H, Baio A, Cassama E, Nabicassa M, Bailey R, et al. Health beliefs and perceptions of trachoma in communities on the Bijagos archipelago of Guinea Bissau. 2015;22(3):190-9.
- 31. Trujillo-Trujillo J, Meza-Cárdenas M, Sánchez SB, Zamora SM, Porras A, López de Mesa CB, et al. Knowledge, Attitudes, and Practices of Hygiene and the Prevention of Trachoma in the Indigenous Population of the Colombian Amazon Vaupés Department. 2023;20(5):4632.
- 32. Yirdaw G, Tegegne EJEHI. Knowledge, Practice and Associated Factor of Trachoma Transmission and Prevention Among Primary School Students in Addis Zemen Town, South Gondar, Northwest Ethiopia. 2024;18:11786302241306933.
- 33. Munguti PN. Knowledge, Practices and Perceptionss of Trachoma and its' Influence (Association With) on Health Seeking Behaviour of Patients in Kajiado Central Sub County, Kajiado County, Kenya: COHES, JKUAT; 2017.
- 34. Solomon AW, Organization WH, Initiative IT. Trachoma control: a guide for programme managers: World Health Organization; 2006.
- 35. Burton MJ, Mabey DCJPntd. The global burden of trachoma: a review. 2009;3(10):e460.
- Hu VH, Harding-Esch EM, Burton MJ, Bailey RL, Kadimpeul J, Mabey DCJTM, et al. Epidemiology and control of trachoma: systematic review. 2010;15(6):673-91.
- Mohammadpour M, Abrishami M, Masoumi A, Hashemi HJJoco. Trachoma: Past, present and future. 2016;28(4):165-9.
- Solomon AW, Burton MJ, Gower EW, Harding-Esch EM, Oldenburg CE, Taylor HR, et al. Trachoma. 2022;8(1):32.
- 39. Stocks ME, Ogden S, Haddad D, Addiss DG, McGuire C, Freeman MCJPm. Effect of water, sanitation, and hygiene on the prevention of trachoma: a systematic review and meta-analysis. 2014;11(2):e1001605.
- 40. Migchelsen SJ. Chlamydia trachomatis serology as a means of monitoring intervention activities to eliminate trachoma as a public health problem: London School of Hygiene & Tropical Medicine; 2019.
- Mtuy T. Maasai Response to Mass Drug Administration for Trachoma in a Changing Political Economy in Tanzania: London School of Hygiene & Tropical Medicine; 2023.
- 42. Alada JJ, Mpyet C, Florea VV, Boisson S, Willis R, Bakhtiari A, et al. Prevalence of trachoma in Kogi State, Nigeria: results of four local government area-level surveys from the global trachoma mapping project. 2018;25(sup1):33-40.
- 43. Cassivi A. Access to drinking water in low-and middle-income countries: monitoring and assessment 2020.
- 44. Adane B, Malede A, Sewunet B, Kumlachew L, Moges M, Woretaw L, et al. Determinants of Trachomatous Inflammation-Follicular Among Children Aged 1 to 9 Years Old in a Rural Area of Gozamn District, Northwestern Ethiopia: A Matched Case-Control Study. 2023;17:11786302231169941.
- 45. LUVISIA MM. INTESTINAL GEOHELMINTH NEMA-TODES INFECTIONS IN LANGAS 2016.
- 46. Tilahun MM, Eticha BL, Shobiso MG, Lorato MM. Knowledge and associated factors towards trachoma in rural Lemo district, Southern Ethiopia. 2024.

- Okumu JO, Gachohi J, Wanjihia VJAJoHS. Water, Sanitation and Hygiene Indicator Levels Eight Years Post Community-Led Total Sanitation Implementation in Kajiado County, Kenya. 2022;35(2):224-40.
- Lakew S, Asefa G, Zerdo ZJBPH. Assessment of the status of improved F&E trachoma control practices among children of agro-pastoralists in Southern Ethiopia: a mixed design survey using theory of triadic influences. 2023;23(1):556.
- 49. Okullo JO, Moturi WN, Ogendi GMJEhi. Open defaecation and its effects on the bacteriological quality of drinking water sources in Isiolo County, Kenya. 2017;11:1178630217735539.
- 50. Bazaanah P. Evaluating the effects of household's socio-demographic elements on the determination of drinking water quality and quantity in rural Savannah communities of Northern Ghana. 2019.

- 51. Damissie F, Aredo MT, Eshete A, Tejineh S, Batu DJS-JoPH. Assessment of Food Safety Practices and Associated Factors Among Health Extension Model and Non-Model Households in Asella Town, South West Ethiopia, 2023. 2024;12(2):103-21.
- 52. Anderson KA. Concerns of Water Scarcity and Water Quality among two Andean Communities in Peru: University of South Florida; 2016.
- Aya Pastrana N, Lazo-Porras M, Miranda JJ, Beran D, Suggs LSJPntd. Social marketing interventions for the prevention and control of neglected tropical diseases: A systematic review. 2020;14(6):e0008360.
- 54. Branda F, Ali AY, Ceccarelli G, Albanese M, Binetti E, Giovanetti M, et al. Assessing the Burden of Neglected Tropical Diseases in Low-Income Communities: Challenges and Solutions. 2024;17(1):29.

Copyright: © 2025 by the contributors and Crystal Scientific Publishers. This work is licensed under the Creative Commons Attribution International License.